

Drug Facts**Active ingredient** **Purpose**
(in each 7.5 mL)Loperamide HCl
1 mg.....Anti-diarrheal**Use** controls symptoms of diarrhea,
including Travelers' Diarrhea**Warnings****Allergy alert:** Do not use if you have
ever had a rash or other allergic
reaction to loperamide HCl**Do not use** if you have bloody or
black stool**PEEL BACK HERE****MAJOR**® NDC 0904-6256-20**Loperamide
Hydrochloride
Oral Suspension**

Anti-Diarrheal Liquid

Controls the
Symptoms of Diarrhea

Mint Flavor

4 FL OZ (120 mL)

1 mg Loperamide Hydrochloride per 7.5 mL

**COMPARE to the
active ingredient of
IMODIUM® A-D*****Do not use if printed
plastic neckband is
broken or missing***Major® Loperamide Hydrochloride Oral Suspension is
not manufactured or distributed by McKel Consumer
Healthcare, distributor of Imodium® A-D.**Distributed by: MAJOR® PHARMACEUTICALS**
31778 Enterprise Drive, Livonia, MI 48150 USA
M-06 REV. 09/13
Re-Order No. 700606**Gluten Free**5
09046 25620
3**CODE
AREA**• no varnish • no color
: 64526 5C F2**ADHESIVE
AREA
NO COPY****Drug Facts** (continued)children 6-8 years
(48-59 lbs)15 mL (3 tsp) after the first
loose stool; 7.5 mL (1 1/2 tsp)
after each subsequent loose
stool; but no more than 30 mL
(6 tsp) in 24 hourschildren under
6 years
(up to 47 lbs)

ask a doctor

Other information

- each 30 mL (6 tsp) contains: sodium 15 mg
- store between 20-25°C (68-77°F)

Drug Facts (continued)

- see side panel for lot number and
expiration date

Inactive ingredients anhydrous citric
acid, carboxymethylcellulose sodium, D&C
yellow no. 10, FD&C blue no. 1, glycerin,
microcrystalline cellulose, natural and artificial
mint flavor, propylene glycol, purified water,
simethicone, sodium benzoate, sucralose,
titanium dioxide, xanthan gum**Questions or comments?**

1-800-719-9260

**ADHESIVE
AREA
NO COPY****Drug Facts** (continued)**Ask a doctor before use if you have** ■ fever
■ mucus in the stool ■ a history of liver disease**Ask a doctor or pharmacist before use if you
are taking antibiotics****When using this product** ■ tiredness,
drowsiness or dizziness may occur. Be careful
when driving or operating machinery.**Stop use and ask a doctor if**

- symptoms get worse
 - diarrhea lasts for more than 2 days
 - you get abdominal swelling or bulging.
- These may be signs of a serious condition.

If pregnant or breast-feeding, ask a health
professional before use.**Keep out of reach of children.** In case of
overdose, get medical help or contact a Poison**Drug Facts** (continued)

Control Center right away. (1-800-222-1222)

Directions ■ drink plenty of clear fluids to
help prevent dehydration caused by diarrhea

- find right dose on chart. If possible, use weight
to dose; otherwise use age.
- shake well
before using
- only use attached measuring
cup to dose product

adults and children
12 years and over30 mL (6 tsp) after the first loose
stool; 15 mL (3 tsp) after each
subsequent loose stool; but no more
than 60 mL (12 tsp) in 24 hourschildren 9-11 years
(60-95 lbs)15 mL (3 tsp) after the first loose
stool; 7.5 mL (1 1/2 tsp) after each
subsequent loose stool; but no more
than 45 mL (9 tsp) in 24 hours