Active ingredients (in each TSP) Dextromethorphan HBr 10 mg Guaifenesin 100 mg

Warnings Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for two weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOL ask a doctor or pharmacist before use.

Ask a doctor before use if you have

 a cough with too much phleam (mucus). a persistent or chronic cough as occurs with smoking, asthma, chronic bronchitis or emphysema. Stop use and ask a doctor if cough lasts more than 7 days, comes back or occurs with fever, rash or headache that lasts. These could be signs of a serious condition. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.



NDC 0536-0970-97

- COLIGH
- NON-NARCOTIC

SYRUP

Guaifenesin with Dextromethorphan HBr

Distributed by: Rugby Laboratories 31778 Enterprise Drive Livonia, MI 48150 4 fl oz (118 mL)

Directions • take every 4 hours use only with enclosed dose cup. Do not use with any other device.

Adults & children 12 years & over | 2 TSP Children under 12 years do not use

TAMPER EVIDENT: DO NOT USE IF

INNER FOIL SEAL OVER MOUTH OF

BOTTLE IS BROKEN OR MISSING

do not take more than 6 doses in 24 hours.

SEE BOX FOR COMPLETE DRUG FACTS

- · each teaspoon contains: sodium 3 mg
- store at 20°-25°C (68°-77°F). Questions or comments? 1-800-645-2158

Re-order No. 370057 R-64

Rev 03/13