



 **MAJOR®**
Original Prescription Strength

FEXOFENADINE HYDROCHLORIDE

Tablets, **180 mg** / Antihistamine

ALLERGY

**24
HOUR**

COMPARE TO the active ingredient of **ALLEGRA® ALLERGY TABLETS***

NDC 0904-6214-46

 **MAJOR®**

Original Prescription Strength

FEXOFENADINE HYDROCHLORIDE

Tablets, **180 mg** / Antihistamine

ALLERGY

Relief of:

- Sneezing
- Runny Nose
- Itchy, Watery Eyes
- Itchy Nose or Throat

Indoor & Outdoor
Allergies



actual size

**24
HOUR**

**30 TABLETS
180 mg EACH**

Non-Drowsy



3 09046 21446 5

Drug Facts (continued)

Inactive ingredients colloidal silicon dioxide, croscarmellose sodium, hypromellose, iron oxide black, iron oxide red, iron oxide yellow, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, titanium dioxide


Questions or comments?
1-800-719-9260

Gluten Free

*Major® Fexofenadine Hydrochloride Tablets, 180 mg are not manufactured or distributed by Chattem, Inc., distributor of Allegra® Allergy Tablets.

Made in Israel

Distributed by
MAJOR®
PHARMACEUTICALS
31778 Enterprise Drive
Livonia, MI 48150 USA
M-05 REV. 09/13
Re-Order No. 265252



Drug Facts

Active ingredient (in each tablet)

Fexofenadine HCl 180 mg.....Antihistamine

Purpose

Uses

temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:

■ runny nose ■ itchy, watery eyes ■ sneezing ■ itching of the nose or throat

Warnings

Do not use if you have ever had an allergic reaction to this product or any of its ingredients.

Ask a doctor before use if you have kidney disease. Your doctor should determine if you need a different dose.

When using this product

- do not take more than directed
- do not take at the same time as aluminum or magnesium antacids
- do not take with fruit juices (see Directions)

Stop use and ask a doctor if an allergic reaction to this product occurs. Seek medical help right away.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

adults and children 12 years of age and over	take one 180 mg tablet with water once a day; do not take more than 1 tablet in 24 hours
children under 12 years of age	do not use
adults 65 years of age and older	ask a doctor
consumers with kidney disease	ask a doctor

Other information

- do not use if printed foil under cap is broken or missing
- store at 20°-25°C (68°-77°F)
- protect from excessive moisture
- this product meets the requirements of USP Dissolution Test 3

