

B-9811R2

Drug Facts

KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION

Active ingredients (in each tablet) Purpose

Pseudoephedrine HCl 60 mg..... Nasal decongestant
Triprolidine HCl 2.5 mg..... Antihistamine

Uses

- temporarily relieves these symptoms due to hay fever (allergic rhinitis) or other upper respiratory allergies:
 - runny nose
 - itchy, watery eyes
 - nasal congestion
 - sneezing
 - itching of the nose or throat
- temporarily relieves these symptoms due to the common cold:
 - runny nose
 - sneezing
 - nasal congestion

Warnings

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

- Ask a doctor before use if you have**
- high blood pressure
 - heart disease
 - thyroid disease
 - diabetes
 - glaucoma
 - a breathing problem such as emphysema or chronic bronchitis
 - difficulty in urination due to enlargement of the prostate gland

Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers.

Drug Facts (continued)

When using this product

- **do not exceed recommended dosage**
- excitability may occur, especially in children
- use caution when driving a motor vehicle or operating machinery
- drowsiness may occur
- avoid alcoholic beverages
- alcohol, sedatives and tranquilizers may increase drowsiness

Stop use and ask a doctor if

- nervousness, dizziness or sleeplessness occur
- symptoms do not improve within 7 days or occur with a fever

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- adults and children 12 years and over: take 1 tablet every 4 to 6 hours. Do not take more than 4 tablets in 24 hours.
- children under 12 years: do not use

Other information

- **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

Drug Facts (continued)

Inactive ingredients colloidal silicon dioxide, corn starch, hypromellose, lactose anhydrous, magnesium stearate, microcrystalline cellulose, polyethylene glycol, stearic acid, titanium dioxide

Drug Facts (continued)

Questions or comments? (800) 616-2471

50844

REV0719N17808

B-1212-178-08-R
REV0719N17808

24 Tablets



Nasal Decongestant/Antihistamine
Sneezing, Runny Nose,
Itchy, Watery Eyes

60 mg/2.5 mg

MAJOR®
Maximum Strength™
Aprodine™ Tablets
Pseudoephedrine HCl 60 mg
Triprolidine HCl 2.5 mg

NDC 0904-0250-24

No print/No varnish
Lot & Exp date

TAMPER EVIDENT: DO NOT USE IF
PACKAGE IS OPENED OR IF BLISTER
UNIT IS TORN, BROKEN OR SHOWS
ANY SIGNS OF TAMPERING



Distributed by:
MAJOR® PHARMACEUTICALS
17177 N Laurel Park Drive, Suite 233
Livonia, MI 48152
Rev. 11/19 M-17
Re-order No. 700796

MAJOR®
Maximum Strength™
Aprodine™ Tablets
Pseudoephedrine HCl 60 mg
Triprolidine HCl 2.5 mg