**MAJOR**°

NDC 0904-0734-31

Compare to the active ingredients in Neosporin® Original Ointment\*

# Original Strength

**Triple Antibiotic Ointment** 

Bacitracin Zinc | Neomycin Sulfate | Polymyxin B Sulfate

#### **First Aid Antibiotic**

For external use only

First aid antibiotic to help prevent infection in minor cuts, scrapes, and burns

NET WT. 1 OZ (28 g)

#### **MAJOR**<sup>®</sup>

**Original Strength** 

### **Triple Antibiotic Ointment**

Bacitracin Zinc | Neomycin Sulfate | Polymyxin B Sulfate

**First Aid Antibiotic** 

# **MAJOR**<sup>®</sup>

**Original Strength** 

## **Triple Antibiotic Ointment**

Bacitracin Zinc | Neomycin Sulfate | Polymyxin B Sulfate

**First Aid Antibiotic** 

For external use only

First aid antibiotic to help prevent infection in minor cuts, scrapes, and burns

NDC 0904-0734-31

Compare to the active ingredients in Neosporin® Original Ointment\*

NET WT. 1 OZ (28 g)

=MBOSSING AREA



Distributed by: MAJOR® PHARMACEUTICALS Livonia, MI 48152

Product of India

Code: GO/DRUGS/362 ABA3B/02

PLD-8693B FC007300
Rev.11/20 M-58 Re-order No.100508

Original Strength
Triple Antibiotic
Ointment
Bacitracin Zinc
Neomycin Sulfate
Polymyxin Sulfate
First Aid Antibiotic

\*This product is not manufactured or distributed by Johnson & Johnson Consumer, Inc., distributor of Neosporin® Original Dintment.

Purpose

८आ (८००) ९१९-५४४। **(nestions or comments**?

Inactive ingredient white petrolatum

Other information

stock et 15°-30°C (59°-86°F). Protect from freezing.

before using any medication, read all label directions. Keep
carton, it contains important information.

■ mash pe coneted whith a sterile bandage antages area of the tip of a finger) on the area 1 to 3 times daily a clean the streeted area — clean the attacted area

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Drug Facts (continued)

DIRECTIONS

Stop use and ask a doctor if wo you need to use longer than one allergic reaction develops

Ask a doctor before use if you have

Do not use ■ it you sre allergic to any of the ingredients ■ in the eyes ■ over large areas of the body

> Warnings For external use only.

**USGS** First sid to help prevent infection in minor cuts

Bacitracin zinc, USP 400 units Neomycin 3.5 mg Polymyxin B sulfate, USP 5,000 units

Active ingredients (in each gram)

Drug Facts