Suppositories

Questions or comments? 1-800-645-2158

Inactive ingredients corn starch, methylparaben, propylparaben

1919er information 1909 30°C (4°68) To avoid melting

cectally up to 4 times daily

e insert suppository into the rectum adults and children 12 years and over: insert one suppository

downward, unwrapping the suppository insert suppository into the rectum

edistant one suppository from the strip
 esparate the two tabs of the white plastic wrap
 with thumb and forefinger of each hand, grasp the tabs and peel
 downward in presentinger of the suppository.

with an appropriate cleansing with a tissue or a soft cloth before insertion of suppository

Directions• when practical, cleanse the affected area by patting or blotting with an appropriate cleansing wipe

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Stop uses and ask a doctor if • bleeding occurs
• condition worsens or does not improve within 7 days

Ask a doctor or pharmacist before use if you are presently taking a prescription drug for high blood prescure or depression When using this product, do not use more than directed

- easeasto hand ea
 - Ask a doctor before use if you have

Warnings For rectal use only

 relieves the itching, burning and discomfort associated with hemorrhoids

Phenylephrine HGI O.25%......Vasoconstrictor

Uses

• temporarily shrinks hemorrhoidal tissue

səsoqru¶ (voitsoqqus dəsə ni) sinəibərgini əvitəA %7.88 fət bisH bish

Drug Facts

NDC 0536-1389-12

Lot/Exp



Easy-To-Use Packaging

Hemorrhoidal Suppositories

Temporarily shrinks hemorrhoids and provides prompt relief from itching, burning and discomfort

12 RECTAL SUPPOSITORIES

TAMPER EVIDENT: DO NOT USE IF FILM IS TORN OR BROKEN

Distributed by: Rugby Laboratories 31778 Enterprise Drive Livonia, MI 48150

Rev. 03/13 R-64 www.rugbylaboratories.com

