



NDC 0904-6464-20



ROBAFEN®

DM COUGH & CHEST
CONGESTION

COUGH SUPPRESSANT (Dextromethorphan HBr)
EXPECTORANT (Guaifenesin)

PEAK COLD

Relieves:

- Cough
- Mucus

Non-Drowsy


COMPARE TO the active ingredients of ROBITUSSIN® COUGH + CHEST CONGESTION DM*

FOR ADULTS
Ages 12 and Over

4 FL. OZ. (118 mL)



359265CC8



NDC 0904-6464-20



ROBAFEN®

DM COUGH & CHEST
CONGESTION

COUGH SUPPRESSANT (Dextromethorphan HBr)
EXPECTORANT (Guaifenesin)

PEAK COLD

Relieves:

- Cough
- Mucus

Non-Drowsy

COMPARE TO the active ingredients of ROBITUSSIN® COUGH + CHEST CONGESTION DM*

FOR ADULTS
Ages 12 and Over

4 FL. OZ. (118 mL)

DO NOT USE IF PRINTED NECKBAND
IS BROKEN OR MISSING

ROBAFEN®
DM COUGH & CHEST
CONGESTION



Drug Facts

Active ingredients (in each 10 mL) Purposes

Dextromethorphan HBr,
USP 20 mg.....Cough suppressant
Guaifenesin, USP 200 mg.....Expectorant

Uses

- temporarily relieves cough due to minor throat and bronchial irritation as may occur with a cold
- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes

Warnings

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis, or emphysema

Stop use and ask a doctor if cough lasts for more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. A persistent cough may be a sign of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided
- keep dosing cup with product
- mL = milliliter
- this adult product is not intended for use in children under 12 years of age

Drug Facts (continued)

age	dose
adults and children 12 years and over	10 mL every 4 hours
children under 12 years	do not use

Other information

- each 10 mL contains: sodium 6 mg
- store at 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients anhydrous citric acid, FD&C red no. 40, flavor, glycerin, high fructose corn syrup, menthol, propylene glycol, purified water, sodium benzoate, sodium citrate, sucralose

Questions or comments?

1-800-616-2471

Gluten Free

*Major® Robafen® DM is not manufactured or distributed by Wyeth LLC, owner of the registered trademark Robitussin®.

Distributed by
MAJOR® PHARMACEUTICALS
17177 N Laurel Park Drive, Suite 233
Livonia, MI 48152
M-05 REV. 02/20
Re-Order No. 700413

Dosage Cup Included

PARENTS:

Learn about teen medicine abuse

www.StopMedicineAbuse.org



3 09046 46420 4

