

Active ingredients (in each 10 mL):

Brompheniramine maleate, USP 2 mg; Dextromethorphan HBr, USP 10 mg; Phenylephrine HCl, USP 5 mg

Uses: temporarily relieves cough due to minor throat and bronchial irritation occurring with a cold, and nasal congestion due to the common cold, hay fever or other upper respiratory allergies. Temporarily relieves these symptoms due to hay fever (allergic rhinitis): runny nose; sneezing; itchy, watery eyes; itching of the nose or throat. Temporarily restores freer breathing through the nose.

Warnings

Do not use to make a child sleepy or if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have: heart disease; high blood pressure; thyroid disease; diabetes; trouble urinating due to an enlarged prostate gland; glaucoma; cough that occurs with too much phlegm (mucus); a breathing problem such as emphysema or chronic bronchitis; cough that lasts or is chronic such as occurs with smoking, asthma, or emphysema.

Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers.

When using this product: do not use more than directed; may cause marked drowsiness; avoid alcoholic beverages; alcohol, sedatives, and tranquilizers may increase drowsiness. Be careful when driving a motor vehicle or operating machinery; excitability may occur, especially in children.

NDC 0904-6463-20



Children's

DimaphenTM DM

ANTI-HISTAMINE, Brompheniramine Maleate
COUGH SUPPRESSANT, Dextromethorphan HBr
NASAL DECONGESTANT, Phenylephrine HCl

Relieves • Nasal Symptoms
• Stuffy Nose • Runny Nose
• Sneezing
Plus Other Symptoms
• Itchy, Watery eyes
• Cough

GRAPE FLAVOR

Alcohol-Free

6 yrs. and older

4 FL. OZ. (118 mL.)

Stop use and ask a doctor if: you get nervous, dizzy, or sleepless; symptoms do not get better within 7 days or are accompanied by fever; cough lasts for more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. A persistent cough may be a sign of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions: do not take more than 6 doses in any 24-hour period; measure only with dosage cup provided; keep dosage cup with product; mL = milliliter.

age	dose
adults and children 12 years and over	20 mL every 4 hours
children 6 to under 12 years	10 mL every 4 hours
children under 6 years	do not use

Other information: each 10 mL contains: sodium 4 mg. Store at 20°-25°C (68°-77°F).

Inactive ingredients anhydrous citric acid, edetate disodium, FD&C blue no. 1, FD&C red no. 40, flavor, glycerin, propylene glycol, purified water, saccharin sodium, sodium benzoate, sorbitol solution

Questions or comments? 1-800-616-2471

Do not use if printed neckband is broken or missing **Gluten Free**

Distributed by
MAJOR[®] PHARMACEUTICALS
17177 N Laurel Park Drive, Suite 233
Livonia, MI 48152
M-05 REV. 03/20 Re-Order No. 700412

**CODE
AREA**

• no varnish • no color

: 98726 5C F6

09046463207