

MAJOR®

NDC 0904-5769-60

Māpap®

ARTHRITIS PAIN

ACETAMINOPHEN
EXTENDED-RELEASE TABLETS, 650 mg
PAIN RELIEVER/FEVER REDUCER

FOR THE TEMPORARY RELIEF
OF MINOR ARTHRITIS PAIN

READ THE
LABEL

DO NOT USE WITH
OTHER MEDICINES
CONTAINING
ACETAMINOPHEN

100 Caplets*
650 mg. Each

*Capsule-Shaped Tablets



CAPLETS

Push & Turn Cap

Active ingredient (in each caplet)

Acetaminophen 650 mg.....Pain reliever/fever reducer

Purpose

Uses ■ temporarily relieves minor aches and pains due to: ■ minor pain of arthritis ■ muscular aches ■ backache ■ premenstrual and menstrual cramps ■ the common cold ■ headache ■ toothache ■ temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take ■ more than 6 caplets in 24 hours, which is the maximum daily amount ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product **Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash If a skin reaction occurs, stop use and seek medical help right away.

Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are allergic to acetaminophen or any of the inactive ingredients in this product. **Ask a doctor before use if you have liver disease.**

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin. Stop use and ask a doctor if ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ new symptoms occur ■ redness or swelling is present. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children. Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms. **Directions** ■ **do not take more than directed (see overdose warning)**

Adults: ■ take 2 caplets every 8 hours with water ■ swallow whole; do not crush, chew, split or dissolve ■ do not take more than 6 caplets in 24 hours ■ do not use for more than 10 days unless directed by a doctor **Under 18 years of age:** ■ ask a doctor **Other information** ■ store at 20-25°C (68-77°F) ■ **do not use if printed foil under cap is broken or missing** ■ meets the requirements of USP *Dissolution Test 2* **Inactive ingredients** carnauba wax, colloidal silicon dioxide, croscarmellose sodium, hypromellose, magnesium stearate, maltodextrin, microcrystalline cellulose, polyethylene glycol, polysorbate 80, povidone, pregelatinized starch, stearic acid, titanium dioxide

Questions or comments? 1-800-719-9260

Distributed by: MAJOR® PHARMACEUTICALS

17177 N Laurel Park Drive, Suite 233

Livonia, MI 48152

M-05 REV. 03/20 Re-Order No. 100381

U.S. Patent
7,897,172

Gluten
Free

: 54478 5C F9

0969257060

Σ