

A441R

Drug Facts (continued)

sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
 ■ if your child is allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if your child has liver disease.

Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin.

When using this product do not exceed recommended dosage.

Stop use and ask a doctor if

- pain gets worse or lasts more than 5 days
 - fever gets worse or lasts more than 3 days
 - new symptoms occur
 - redness or swelling is present
- These could be signs of a serious condition.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical even if you do not notice any signs or symptoms.

Drug Facts (continued)

Directions ■ **this product does not contain directions or complete warnings for adult use**

- find the right dose on chart below. If possible, use weight to dose; otherwise, use age.
- dissolve in mouth or chew before swallowing
- if needed, repeat dose every 4 hours while symptoms last
- do not give more than 5 times in 24 hours
- do not give for more than 5 days unless directed by a doctor

| Weight (lb) | Age (yr) | Tablets |
|-------------|----------|--------------|
| Under 48 | Under 6 | Ask a doctor |
| 48-59 | 6-8 | 2 |
| 60-71 | 9-10 | 2 1/2 |
| 72-95 | 11 | 3 |

Other information

- **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

Drug Facts (continued)

Inactive ingredients citric acid, crospovidone, D&C red #27 aluminum lake, D&C red #30 aluminum lake, dextrates hydrated, ethylcellulose, flavor, magnesium stearate, mannitol, polyethylene, stearic acid, sucralose

Questions or comments? (800) 616-2471

*This product is not manufactured or distributed by McNeil Consumer Healthcare, owner of the registered trademark Junior Strength Tylenol® Meltaways®. 50844 REV0213E45008



B-1212-450-08
REV0213E45008

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

Bubble Gum Flavor

24 Tablets, 160 mg EACH

AGES 6-11

Rapid Tabs

ASPIRIN FREE • IBUPROFEN FREE

Fever reducer / Pain reliever

ACETAMINOPHEN

Junior Mappap

Compare to the active ingredient in Junior Tylenol® Meltaways®
NDC 0904-5754-24

MAJOR

Distributed by MAJOR® PHARMACEUTICALS
31778 Enterprise Drive
Livonia, MI 48150 USA
M-17
Rev. 05/13
Re-order No. 100178

No Print Area
Lot no. & Exp. Date

Drug Facts

KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION

Active ingredient (in each tablet) Purpose
 Acetaminophen 160 mg... Pain reliever/fever reducer

Uses ■ temporarily relieves minor aches and pains due to: ■ the common cold ■ flu ■ headache ■ sore throat ■ temporarily reduces fever

Warnings
Liver warning: This product contains acetaminophen.

Drug Facts (continued)

Severe liver damage may occur if your child takes:

- more than 5 doses in 24 hours, which is the maximum daily amount
 - with other drugs containing acetaminophen
- Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not