

MAJOR® NDC 0904-0250-59

Maximum Strength

Aprodine™ Tablets

Pseudoephedrine HCl 60 mg

Triprolidine HCl 2.5 mg

60 mg/2.5 mg

Nasal Decongestant
Antihistamine

Relieves Nasal Congestion,
Sneezing, Runny Nose,
Itchy, Watery Eyes

100 Tablets

50844 REV0719K17812

Dist. by: **MAJOR® PHARMACEUTICALS**
17177 N Laurel Park Drive, Suite 233
Livonia, MI 48152

Rev. 11/19 M-17 Re-order No. 700801



**KEEP OUTER PACKAGE FOR
COMPLETE PRODUCT INFORMATION**

Drug Facts

Active ingredients (in each tablet) Purpose

Pseudoephedrine HCl 60 mg..... Nasal decongestant

Triprolidine HCl 2.5 mg..... Antihistamine

Uses ■ temporarily relieves these symptoms due to hay

fever (allergic rhinitis) or other upper respiratory allergies:

■ runny nose ■ itchy, watery eyes ■ nasal congestion

■ sneezing ■ itching of the nose or throat

■ temporarily relieves these symptoms due to the common

cold: ■ sneezing ■ nasal congestion ■ runny nose

Warnings See carton for complete warnings

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

■ adults and children 12 years and over: take 1 tablet every

4 to 6 hours. Do not take more than 4 tablets in 24 hours.

■ children under 12 years: do not use

Other information

■ **TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY**

SEAL UNDER CAP IS BROKEN OR MISSING

■ use by expiration date on package

■ store at 25°C (77°F); excursions permitted between

15°-30°C (59°-86°F)