



**NEW!**

# KLENTZ™ (florfenicol, terbinafine, mometasone furoate) Otic Solution

## TWIST-N-APPLY

tube design that simplifies and enables precise application to the desired area. Enjoy a targeted and gentle application with the **soft, flexible nozzle design**

Approved by FDA under ANADA #200-829

## Featuring:

- 16.6 mg/ml florfenicol, 14.8 mg/ml terbinafine (equivalent to 16.6 mg/ml of terbinafine hydrochloride) and 2.2 mg/ml mometasone furoate
- 1-dose dropperette for all canine sizes
- Single dose treatment regimen
- A non-viscous liquid solution

**Cost-effective,  
bioequivalent to Claro™**



TWIST-N-APPLY



## AVAILABLE PACKS

- Reorder no: 16030 - **2 pk**
- Reorder no: 16031 - **12 pk**
- Reorder no: 16032 - **24 pk**

**Unique, flexible  
soft tip applicator**



### IMPORTANT SAFETY INFORMATION:

**Do not use in cats. KLENTZ™ should be administered by veterinary personnel.** KLENTZ™ may cause eye injury and irritation. Splatter may occur if the dog shakes its head following administration. If contact with eyes occurs, flush copiously with water for at least 15 minutes. If irritation persists, contact a physician. Do not use in dogs with known tympanic membrane perforation (ruptured eardrum). Adverse reactions may occur following administration of KLENTZ™. Observe your dog for signs such as ear pain, irritation, vomiting, head shaking, head tilt, in-coordination, eye pain and discharge. Contact your veterinarian if any of these signs are observed. For complete safety information, please see KLENTZ™ package insert on the reverse side.

# Klantz™

## (florfenicol, terbinafine, mometasone furoate)

### Otic Solution for use in dogs only

#### Do Not Use in Cats.

Antibacterial, antifungal, and anti-inflammatory

**CAUTION:** Federal law restricts this drug to use by or on the order of a licensed veterinarian.

#### DESCRIPTION:

Klantz contains 16.6 mg/mL florfenicol, 14.8 mg/mL terbinafine (equivalent to 16.6 mg/mL terbinafine hydrochloride) and 2.2 mg/mL mometasone furoate. Inactive ingredients include purified water, propylene carbonate, propylene glycol, ethyl alcohol, and polyethylene glycol.

#### INDICATIONS:

Klantz is indicated for the treatment of otitis externa in dogs associated with susceptible strains of yeast (*Malassezia pachydermatis*) and bacteria (*Staphylococcus pseudintermedius*).

#### DOSAGE AND ADMINISTRATION:

**Klantz should be administered by veterinary personnel.**

**Wear eye protection when administering Klantz.** (see **Human Warnings, Precautions, Post-Approval Experience**).




Splatter may occur if the dog shakes its head following administration. Persons near the dog during administration should also take steps to avoid ocular exposure.

**Shake before use.**

**Verify the tympanic membrane is intact prior to administration.** (see **Contraindications, Precautions, Post-Approval Experience**).

Administer one dose (1 dropperette) per affected ear.

1. Clean and dry the external ear canal before administering the product.
2. Verify the tympanic membrane is intact prior to administration.
3. Remove single dose dropperette from the package.
4. While holding the dropperette in an upright position, twist the cap to open the dropperette. The cap does not come off the dropperette.



5. Insert the tapered tip of the dropperette into the affected external ear canal and squeeze to instill the entire contents (1 mL) into the affected ear.
6. Gently massage the base of the ear to allow distribution of the solution. **Restrain the dog to minimize post application head shaking** to reduce potential for splatter of product and accidental eye exposure in people and dogs (see **Post-Approval Experience**).



7. Repeat with other ear as prescribed.
8. The duration of the effect should last 30 days. Cleaning the ear after dosing may affect product effectiveness.

#### CONTRAINDICATIONS:

Do not use in dogs with known tympanic membrane perforation (see **Precautions**).

Klantz is contraindicated in dogs with known or suspected hypersensitivity to florfenicol, terbinafine hydrochloride, or mometasone furoate.

#### WARNINGS:

**Human Warnings:** Klantz may cause eye injury and irritation (see **Precautions, Post-Approval Experience**).

If contact with eyes occurs, flush copiously with water for at least 15 minutes. If irritation persists, contact a physician.

Humans with known hypersensitivity to any of the active ingredients in Klantz should not handle this product.

Not for use in humans. Keep this and all drugs out of reach of children. Avoid skin contact. In case of accidental ingestion by humans, contact a physician immediately.

#### PRECAUTIONS:

**For use in dogs only. Do not use in cats** (see **Post-Approval Experience**).

**Wear eye protection when administering Klantz and restrain the dog** to minimize post application head shaking. Reducing the potential for splatter of product will help prevent accidental eye exposure in people and dogs and help to prevent ocular injury (see **Dosage and Administration, Human Warnings, Post-Approval Experience**).

Proper patient selection is important when considering the benefits and risks of using Klantz. The integrity of the tympanic membrane should be confirmed before administering the product.

Florfenicol, terbinafine, and mometasone furoate otic solution has been associated with rupture of the tympanic membrane. Reevaluate the dog if hearing loss or signs of vestibular dysfunction are observed during treatment.

Signs of internal ear disease such as head tilt, vestibular signs, ataxia, nystagmus, facial paralysis, and keratoconjunctivitis sicca have been reported (see **Post-Approval Experience**) with the use of florfenicol, terbinafine, and mometasone furoate otic solution.

Do not administer orally.

Use of topical otic corticosteroids has been associated with adrenocortical suppression and iatrogenic hyperadrenocorticism in dogs (see **Animal Safety**).

Use with caution in dogs with impaired hepatic function (see **Animal Safety**).

The safe use of Klantz in dogs used for breeding purposes, during pregnancy, or in lactating bitches, has not been evaluated.

#### ADVERSE REACTIONS:

In a field study conducted in the United States (see **Effectiveness**), there were no directly attributable adverse reactions in 146 dogs administered florfenicol, terbinafine, and mometasone furoate otic solution.

#### POST-APPROVAL EXPERIENCE (2019):

The following adverse events are based on post-approval adverse drug experience reporting for florfenicol, terbinafine, and mometasone furoate otic solution. Not all adverse events are reported to FDA/CVM. It is not always possible to reliably estimate the adverse event frequency or establish a causal relationship to product exposure using these data.

In **humans**, accidental exposure leading to corneal ulcers and other ocular injuries such as eye irritation and redness have been reported. Exposure occurred when the dog shook its head after application of florfenicol, terbinafine, and mometasone furoate otic solution.

Skin irritation has also been reported.

In **dogs**, the adverse events reported are presented below in decreasing order of reporting frequency:

Ear discharge, head shaking, ataxia, internal ear disorder (head tilt and vestibular), deafness, emesis, nystagmus, pinnal irritation and ear pain, keratoconjunctivitis sicca, vocalization, corneal ulcer, cranial nerve disorder (facial paralysis), tympanic membrane rupture.

Klantz (florfenicol, terbinafine, mometasone furoate) otic solution is not approved for use in **cats**. The adverse events reported following extra-label use in **cats** are presented below in decreasing order of reporting frequency: Ataxia, anorexia, internal ear disorder (head tilt and vestibular), Horner's syndrome (third eyelid prolapse and miosis), nystagmus, lethargy, anisocoria, head shake, emesis, tympanic rupture, and deafness.

#### CONTACT INFORMATION:

To report suspected adverse drug experiences, for technical assistance, or to obtain a copy of the Safety Data Sheet (SDS), contact Aurora Pharmaceutical at 1-888-215-1256.

For additional information about reporting adverse drug experiences for animal drugs, contact FDA at 1-888-FDA-VETS or <http://www.fda.gov/reportanimalae>

#### Information for Dog Owners:

Owners should be aware that adverse reactions may occur following administration of Klantz and should be instructed to observe the dog for signs such as ear pain and irritation, vomiting, head shaking, head tilt, incoordination, eye pain and ocular discharge (see **Post-Approval Experience**). Owners should be advised to contact their veterinarian if any of the above signs are observed.

Owners should also be informed that splatter may occur if the dog shakes its head following administration of Klantz which may lead to ocular exposure. Eye injuries, including corneal ulcers, have been reported in humans and dogs associated with head shaking and splatter following administration. Owners should be careful to avoid ocular exposure (see **Precautions, Post-Approval Experience**).

#### PHARMACOLOGY:

Klantz otic solution is a fixed combination of three active substances: florfenicol (antibacterial), terbinafine (antifungal), and mometasone furoate (steroidal anti-inflammatory). Florfenicol is a bacteriostatic antibiotic which acts by inhibiting protein synthesis. Terbinafine is an antifungal which selectively inhibits the early synthesis of ergosterol. Mometasone furoate is a glucocorticosteroid with anti-inflammatory activity.

#### MICROBIOLOGY:

The compatibility and additive effect of each of the components in florfenicol, terbinafine, and mometasone furoate otic solution was demonstrated in a component effectiveness and non-interference study. An *in vitro* study of organisms collected from clinical cases of otitis externa in dogs enrolled in the clinical effectiveness study determined that florfenicol and terbinafine hydrochloride inhibit the growth of bacteria and yeast commonly associated with otitis externa in dogs. No consistent synergistic or antagonistic effect of the two antimicrobials was demonstrated. The addition of mometasone furoate to the combination did not impair antimicrobial activity to any clinically significant extent. In a field study (see **Effectiveness**), at least 10 isolates from successfully treated cases were obtained for *S. pseudintermedius* and *M. pachydermatis*.

#### EFFECTIVENESS:

In a well-controlled, double-masked field study, florfenicol, terbinafine, and mometasone furoate otic solution was evaluated against a vehicle control in 221 dogs with otitis externa. One hundred and forty six dogs were treated with florfenicol, terbinafine, and mometasone furoate otic solution and 75 dogs were treated with the vehicle control. All dogs were evaluated for safety. Treatment (1 mL) was administered once on Day 0 to the affected ear(s). Prior to treatment, the ear(s) was cleaned with saline. The dogs were evaluated on Days 0, 7, 14, and 30. Blood work and urinalysis were obtained on Day 0 pre-treatment and Day 30 at study completion. Four clinical signs associated with otitis externa were evaluated: erythema, exudate, swelling, and ulceration. Success was based on clinical improvement at Day 30. Of the 183 dogs included in the effectiveness evaluation, 72.5% of dogs administered florfenicol, terbinafine, and mometasone furoate otic solution were successfully treated, compared to 11.1% of the dogs in the vehicle-control group ( $p = 0.0001$ ).

#### ANIMAL SAFETY:

In a target animal safety study, florfenicol, terbinafine, and mometasone furoate otic solution was administered aurally to 12-week-old Beagle puppies (4 dogs/sex/group) at 0X, 1X, 3X, and 5X the recommended dose once every 2 weeks for a total dosing period of 28 days (3 times the treatment duration). No clinically relevant treatment-related findings were noted in hearing tests, body weight, weight gain, or food consumption. Florfenicol, terbinafine, and mometasone furoate otic solution administration was associated with post-treatment ear wetness or clear aural exudate, increased absolute neutrophil count, decreased absolute lymphocyte and eosinophil counts, suppression of the adrenal cortical response to ACTH-stimulation, decreased adrenal weight and atrophy of the adrenal cortex, increased liver weight with hepatocellular enlargement/cytoplasmic change, and decreased thymus weight. Other potentially treatment-related effects included mild changes to AST, total protein, inorganic phosphorus, creatinine, and calcium.

#### STORAGE INFORMATION:

Store up to 25°C (77°F), excursions are permitted up to 30°C (86°F).

#### HOW SUPPLIED:

Klantz solution is supplied in a single-use dropperette in a blister.

Each dropperette contains one 1 mL dose.

Klantz is available in cartons of two, twelve, or twenty-four dropperettes.

Manufactured by:

Aurora Pharmaceutical, Inc.  
Northfield, MN 55057

1-888-215-1256 [www.aurorapharmaceutical.com](http://www.aurorapharmaceutical.com)

Approved by FDA under ANADA # 200-829

Rev No. 11/2025

